

# Complex Spectral Variability from Intensive Multi-wavelength Monitoring of Mrk 421 in 1998

T. Takahashi<sup>1,2</sup>, J. Kataoka<sup>1,2</sup>, G. Madejski<sup>3,4</sup>, J. Mattox<sup>5</sup>, C.M. Urry<sup>6</sup>, S. Wagner<sup>7</sup>, F. Aharonian<sup>8</sup>, M. Catanese<sup>9</sup>, L. Chiappetti<sup>10</sup>, P. Coppi<sup>11</sup>, B. Degrange<sup>12</sup>, G. Fossati<sup>13</sup>, H. Kubo<sup>14</sup>, H. Krawczynski<sup>8</sup>, F. Makino<sup>1</sup>, H. Marshall<sup>15</sup>, L. Maraschi<sup>16</sup>, F. Piron<sup>12</sup>, R. Remillard<sup>15</sup>, F. Takahara<sup>17</sup>, M. Tashiro<sup>2</sup>, H. Terasranta<sup>18</sup>, and T. Weekes<sup>9</sup>

## ABSTRACT

We conducted a multi-frequency campaign for the TeV blazar Mrk 421 in 1998 April. The campaign started from a pronounced high amplitude flare recorded by *SAX* and *Whipple*; *ASCA* observation started three days later. In the X-ray data, we detected multiple flares, occurring on time scales of about one day. *ASCA* data clearly reveal spectral variability. The comparison of the data from *ASCA*, *EUVE* and *RXTE* indicates that the variability amplitudes in the low energy synchrotron component

---

<sup>1</sup>Institute of Space and Astronautical Science, Kanagawa 229-8510, Japan

<sup>2</sup>Department of Physics, University of Tokyo, Tokyo 113-0033, Japan

<sup>3</sup>Laboratory for High Energy Astrophysics, NASA/GSFC, Greenbelt, MD 20771

<sup>4</sup>Department of Astronomy, University of Maryland, College Park, MD 20742

<sup>5</sup>Astronomy Department, Boston University, Boston, MA, 02215

<sup>6</sup>Space Telescope Science Institute, Baltimore, MD 21218

<sup>7</sup>Landessternwarte Heidelberg, Königstuhl, D-69117 Heidelberg, Germany

<sup>8</sup>Max-Planck-Institut Für Kernphysic, D-69029 Heidelberg, Germany

<sup>9</sup>Fred Lawrence Whipple Observatory, Harvard-Smithsonian CfA, Amado, AZ 85645

<sup>10</sup>Istituto di Fisica Cosmica CNR, Milano, Italy

<sup>11</sup>Department of Astronomy, Yale University, New Haven, CT 06520-8101

<sup>12</sup>LPNHE Ecole Polytechnique Palaiseau, France

<sup>13</sup>University of California at San Diego, La Jolla, CA 92093-0424

<sup>14</sup>Department of Physics, Tokyo Institute of Technology, Tokyo, 152-8551, Japan

<sup>15</sup>Center for Space Research, Cambridge, MA 02139

<sup>16</sup>Osservatorio Astronomico di Brera, 20121 Milano, Italy.

<sup>17</sup>Department of Earth and Space Science, Osaka University, Osaka, Japan

<sup>18</sup>Metsähove Radio Observatory, Kylmala, Finland

are larger at higher photon energies. In TeV  $\gamma$ -rays, large intra-day variations – which were correlated with the X-ray flux – were observed when results from three Cherenkov telescopes are combined. The RMS variability of TeV  $\gamma$ -rays was similar to that observed in hard X-rays, above 10 keV. The X-ray light curve reveals flares which are almost symmetric for most of cases, implying the dominant time scale is the light crossing time through the emitting region. The structure function analysis based on the continuous X-ray light curve of seven days indicates that the characteristic time scale is  $\sim 0.5$  day. The analysis of ASCA light curves in various energy bands appears to show both soft (positive) and hard (negative) lags. These may not be real, as systematic effects could also produce these lags, which are all much smaller than an orbit. If the lags of both signs are real, these imply that the particle acceleration and X-ray cooling time scales are similar.

*Subject headings:* BL Lacertae objects: general – BL Lacertae objects: individual (Markarian 421) – X-rays: general

## 1. Introduction

According to well-established unified schemes for radio-loud AGN (Urry & Padovani 1995), blazars are distinguished by the fortuitous alignment of their jets along the line of sight. This greatly enhances the jet radiation, making them ideal sources for understanding jet physics in all radio-loud AGN. The required high energies of the radiating electrons indicate very efficient particle acceleration in the relativistic jets of these sources (Kubo et al. 1998; Ghisellini et al. 1998).

The BL Lac object Mrk 421 is unique as the first – and so far, the brightest – High Frequency Peaked BL Lac objects (HBLs) with  $\gamma$ -ray emission extending up to TeV energies, at a level allowing detailed spectral and variability studies in the broadest spectral range. The simple continuum spectra of Mrk 421 from radio to UV and X-ray bands obtained previously imply that the emission in these energy bands is due to charged particles radiating via the synchrotron process (e.g., George, Warwick, & Bromage 1988). The hard X-rays and  $\gamma$ -rays from this object are likely to be produced by inverse Compton scattering of synchrotron photons by the same electrons (e.g., Ulrich, Maraschi, & Urry 1997 and references therein). Judging from the results of previous TeV and multi-wavelength monitoring campaigns of Mrk 421, correlations among light curves at different energies (and, in particular, the pronounced spectral evolution during flares seen in X-rays) have provided our best opportunity to understand the high energy emission from blazar jets (Macomb et al. 1995; Buckley et al. 1996; Takahashi et al. 1996; Wagner 1996; Takahashi et al. 1999). Specifically, the previous (1994) campaigns found a soft lag of about 4000 s for soft X-ray (0.5 – 1.0 keV) photons with respect to the hard X-ray (2 – 7.5 keV) band, which was interpreted as an effect of radiative cooling (Takahashi et al. 1996). However, the

sparse sampling of previous campaigns has prevented us from obtaining definitive conclusions. Uninterrupted monitoring, lasting several times the characteristic time scale, is the only way to investigate quantitatively the extreme physical conditions at the sites of high-energy radiation.

Accordingly, we carried out a one-week continuous observation with *ASCA*, coordinated with *EUVE*, and *SAX*. At the same time, TeV detectors (CAT, HEGRA, and Whipple), optical telescopes, radio antennae at 22 GHz and 37 GHz attempted to monitor the source every night. In this paper, we only present the most important result of the multi-wavelength campaign.

## 2. Observations and Results

### 2.1. Observations

The essential information on the observations is summarized in Fig. 1, which shows the normalized multi-wavelength (EUV, X-ray and TeV) light curves. The logarithmic flux scale allows flares with equal amplitudes to be represented by equal vertical excursions. Complete details about the data analyses will be given in separate wavelength-based papers (Aharonian et al. 1999; Fossati et al. 2000a,b; Kataoka et al. 2000a); here we summarize the key points.

Continuous coverage in X-rays with *ASCA* lasted from 1998 April 23.97 to April 30.8 UT, yielding a net exposure of  $\sim 280$  ks. The results presented here are from the SIS detectors operated in 1-CCD mode because the count rate of the GIS detectors exceeded its telemetry limit for 20 – 30 % of the observation time. The background count rate and its fluctuations are negligible, so we did not perform any background subtraction, so as to avoid any artificial effects.

*SAX* observations of Mrk 421 preceded the *ASCA* observation by about three days (Maraschi et al. 1999). *RXTE* observed the source primarily only in low background orbits in order to minimize uncertainties in the background subtraction. *EUVE* observations covered the whole campaign, lasting from April 19.4 to May 1.2. The  $\gamma$ -ray observations at TeV energies were made with the CAT, HEGRA and Whipple atmospheric Cherenkov telescopes. Optical observations with BVRI filters were performed by the WEBT collaboration (Mattox et al., in preparation). Radio observations were done at the Metsähovi Radio Observatory (Teräsranta et al. 1998).

### 2.2. Multi-wavelength light curves

The campaign began during a pronounced high amplitude flare recorded simultaneously by *SAX* and *Whipple* (Maraschi et al. 1999), and the high activity of Mrk 421 continued throughout the campaign (Fig. 1). The 2 – 10 keV flux at the beginning of the *ASCA* observation was  $1.2 \times 10^{-10}$  erg cm $^{-2}$  s $^{-1}$  and increased up to  $5.0 \times 10^{-10}$  erg cm $^{-2}$  s $^{-1}$  at the maximum. Thanks to the continuous coverage with *ASCA*, the X-ray variations are fully resolved *with no gaps*.

Many “flares” are seen clearly in the *ASCA* data, superimposed on a general increasing trend. The *EUVE* light curve also shows a series of flares, less well-defined because of lower amplitude and lower signal-to-noise ratio but well correlated in time with the X-ray flares. The optical data for the period 24 – 30 April indicate an R-band flux with mean value 12.6 mag, and variability of less than 0.1 mag about this mean. The high frequency radio flux did not change within statistics; the mean fluxes during the Metsähovi observations are 0.5 Jy for 22 GHz (20-26 April) and 0.6 Jy for 37 GHz (18-21 April).

In Mrk 421, the EUV through X-ray emission comprises the high energy part of the synchrotron component. Comparison of the data from *ASCA*, *EUVE* and *RXTE* indicates that the variability amplitude in the synchrotron component increases with photon energy. To show this quantitatively, we computed the fractional RMS variability including the general, increasing trend. The calculated RMS variability during *ASCA* campaign increases with energy, from  $0.158 \pm 0.007$  in the *EUVE* light curve, to  $0.195 \pm 0.0004$  in *ASCA* (0.5 – 7 keV), to  $0.385 \pm 0.001$  and  $0.67 \pm 0.03$  for 8 – 16 keV and 16 – 40 keV in the *RXTE* PCA data, respectively. This energy dependence is clear in the multi-epoch multi-frequency spectrum of Mrk 421 shown in Fig. 2, where the largest changes in the synchrotron component occur at the highest frequencies.

The complete flare recorded both in X-rays by *SAX* and in TeV  $\gamma$ -rays by the Whipple telescope demonstrates the correlation between X-ray and TeV  $\gamma$ -ray emission on the time scale of hours (Maraschi et al. 1999). The TeV flux also follows the general week-long rise of the *ASCA* flux (Fig. 10 of Takahashi et al. 1999). Assuming the cross-calibration among the three TeV telescopes is accurate, the relative TeV intensities at nearby times indicate that large intra-day variations are occurring at these high energies. Fractional TeV variability is respectively  $0.30 \pm 0.07$  for the Whipple data and  $0.45 \pm 0.08$  for the HEGRA data. Note that, for the Whipple data,  $F_{var} = 0.43 \pm 0.05$  during the whole campaign. These values are similar to the values for the hard X-ray data above 8 keV, consistent with the scenario that both TeV and hard X-ray emissions are due to the same electron distribution.

In the subsequent discussion, we divided the *ASCA* light curve into segments as marked in Fig. 3, with one flare per segment. In the *ASCA* light curve, most of flares are characterized by nearly equal rise and decay times, i.e., by a nearly symmetric time profile. The time constants from the bottom to the top are also similar from one flare to the next,  $\tau \sim 0.5$  day.

### 2.3. Time series analysis

The long, uninterrupted *ASCA* light curve gives an excellent opportunity to study the temporal characteristics of the X-ray variability. We therefore calculated the first-order *normalized* structure function (SF) at time lag  $\tau$ , which is defined as  $SF(\tau) = \frac{1}{N} \sum [(F(t) - F(t + \tau))/F_{av}]^2$ , where  $F(t)$  is the count rate and  $F_{av}$  is the average count rate.  $N$  is the total number of data pairs used in the calculation. The SF is closely related to the power density spectrum (PDS) distribution

(its slope,  $\beta_{SF}$ , is one less than the PDS slope) but is more suitable when the data sampling is uneven. The SF calculated from the 2 – 7.5 keV *ASCA* light curve is shown in Fig. 4. It has a relatively steep slope at high temporal frequencies (short time scales),  $\beta_{SF} \sim 1.2$  (corresponding to  $\beta_{PDS} \sim 2.2$ ). This is at the steep end of the slopes seen in other types of mass-accreting black hole systems, such as Seyfert galaxies (Hayashida et al. 1998).

A sharp break in the structure function of Mrk 421 occurs at  $\sim 0.5$  days, to a flatter slope at low temporal frequencies (long time scales),  $\beta_{SF} \sim 0.3$  ( $\beta_{PDS} \sim 1.3$ ). This is the first report of a clear turnover in the structure function of a blazar X-ray light curve, and it suggests there is at least one characteristic time scale for the variability of Mrk 421. As shown in Fig. 4, the same flat slope at longer lag is seen in the structure function for the *RXTE* All-Sky Monitor (ASM) light curve (2 – 10 keV), thus confirming the reality of the turnover. In order for the total variability power not to diverge, the structure function must flatten still further at longer time scales; we infer this from the ASM data to be at  $\gtrsim 100$  days. The shape of the structure function is roughly independent of energy from 0.07 – 7.5 keV, as can be seen from the comparison of structure functions for the *ASCA* data and the *EUVE* data.

## 2.4. Spectral Variabilities and Time Lags

The normalized light curves obtained with the SIS in two energy bands (0.5 – 1 keV and 3 – 7.5 keV) are shown in Figs. 3a and 3b, respectively. In general, the amplitude of the variability is larger in hard X-rays than in soft X-rays. As can be seen from the hardness ratio (Fig. 3c), the spectrum becomes harder when the source gets brighter. Generally the spectral changes during individual flares are complex and rapid but in some cases, the hardness ratio remains almost constant. In the flare #8, there is a plateau in the soft X-ray band, which is rarely seen. According to detailed analysis of the spectral variability (Kataoka 2000b), flare #1 shows clockwise hysteresis in the correlation between flux and photon index, similar to that reported by Takahashi et al. (1996a). In contrast, we found anti-clockwise hysteresis in flare #7. In order to quantify the lag, we used the discrete correlation function (DCF; Edelson & Krolik 1988). First, we calculated the DCF for the *ASCA* data comparing each of several energy bands to the 4 – 7.5 keV band using all data. Our results are consistent with zero-lag. We find the same result even using the *EUVE* data, which offers the largest separation in energy from the hard *ASCA* band. However, inspection of X-ray light curves at different energies does suggest some energy dependence of the flare shape. We therefore calculated the lags separately for each flare (as marked in Fig. 3) to see if individual flares have different behavior.

Fig. 3d shows the time lag between 0.5 – 1 keV photons (the softest *ASCA* band) and 4.0 – 7.5 keV photons, calculated for each flare. Errors of lags are determined from the quadratic sum of the uncertainty in determining the maximum of the DCF distribution and the results of MonteCarlo calculation which simulate the effect of time windows (Kataoka 2000b). There are both soft (positive) and hard (negative) lags, with some flares consistent with zero. The soft lag

for flare #1 is  $2000 \pm 710$  s for 0.5 – 1.0 keV photons, which is roughly half that found in previous monitoring in 1994 (Takahashi et al. 1996). In contrast, the lag calculated for flare #7 is negative (a hard lag),  $-3400 \pm 980$  s. These lags may not be real, as it is not possible to reliably determine lags which are much smaller than an orbit. Systematic effects could also produce these lags.

### 3. Discussion

The present multi-wavelength monitoring campaign showed that Mrk 421 is much more variable, and that the variability, and its energy dependence, is more complex than anticipated. The object flares daily (or perhaps even more often), and the correlation of variations in the X-ray and TeV energy bands is confirmed, supporting the idea that the same electron distribution in the same physical region is responsible for the emission in both energy bands. The 2 – 7 keV energy index obtained with *ASCA* ranges from  $\alpha = 1.4$  to 1.8. The energy ranges covered by *ASCA* and *RXTE* are thus at or above the peak in the  $\nu F_\nu$  spectrum. The increasing amplitude of variability at the higher X-ray energies is therefore consistent with the expectations for the synchrotron process, specifically that the higher energy electrons lose their energy more rapidly.

The X-ray variability of Mrk 421 is very similar to the intra-day variability observed in the optical light curves of flat spectrum radio quasars (FSRQs), for which the high-energy component peaks in the GeV  $\gamma$ -ray energy band (Wagner et al. 1996). For synchrotron radiation due to a single population of relativistic electrons with a broken power law distribution of index  $\gamma_{el}$  and a break at  $\gamma_{peak}$ , the correspondign frequency for  $\gamma_{peak}$  is (as measured in the observer’s frame) is  $\nu_{sync} = 3.7 \times 10^6 \gamma_{peak}^2 B \frac{\delta}{(1+z)}$  Hz. Here  $B$  is magnetic field in Gauss, measured in the comoving frame, and  $\delta$  is the Doppler “beaming” factor defined as  $\delta = \Gamma_j^{-1}(1 - \beta \cos \theta)^{-1}$ , where  $\Gamma_j$  is the Lorentz factor of the jet,  $\beta = v/c$ , and  $\theta$  the angle of the jet to the line of sight.

Several analyses have shown that  $\gamma_{peak}$  is lower for FSRQs ( $10^3 - 10^4$ ) and higher for HBLs ( $\sim 10^5 - 10^6$ ), as is the case for Mrk 421 (Sambruna et al. 1996; Kubo et al. 1998; Fossati et al. 1998). Since the synchrotron emission due to the electrons with  $\gamma_{peak}$  corresponds to the eV (or sub-eV) range for FSRQs and the keV range for Mrk 421, the rapid, large-amplitude variability observed above  $\gamma_{peak}$  is likely the result of the rapid change of the electron distribution, in the regime where acceleration and cooling are approximately balanced (e.g., Inoue & Takahara 1996; Kirk et al. 1998). Indeed, the structure function of Mrk 421 derived from the X-ray light curve is very similar to the structure functions found in optical and radio studies of FSRQs, with a break at a characteristic time scale of one day (Wagner & Witzel 1995; Wagner et al. 1996).

The *ASCA* light curve constrains the dominant variability time scales to be less than one day. The quasi-symmetric flare profiles imply that  $\tau_{acc}$  and  $\tau_{cool}$  are shorter than the source light-crossing time  $\tau_{crs}$ , because faster time scales will always be smoothed out by  $\tau_{crs}$  (Chiaberge & Ghisellini 1999; Kataoka et al. 2000c). A plateau also occurs when the time scale for injecting energetic electrons,  $\tau_{inj}$ , is longer than  $\tau_{cool}$ . In this situation, emission from individual slices does

not fade as quickly after an increase, hence the observation of a plateau in the light curve occurs when the entire volume is radiating. We do observe a plateau in the last flare (segment #8), which is much more obvious at low energies, where the cooling time is longer.

A simple conceptual scenario for multiple flares is that a cloud, or a “blob” of plasma passes through the region where shock fronts are formed and electrons are accelerated. If we assume the opening angle of the jet is proportional to  $1/\Gamma_j$ , where the size of the blob at distance  $D$  from the base of the jet can be expressed as  $R \sim D/\Gamma_j$ . With this, the characteristic time scale of  $\tau \sim 0.5$  days (from the structure function analysis) implies that the emission region has size  $R = c\tau\Gamma_j \sim 10^{16}$  cm at distance  $D \sim 10^{17}$  cm from the black hole, if we assume that  $\Gamma_j \sim 10$ . Further discussion of the characteristic time scale and a comparison with other TeV blazars and Seyfert galaxies is presented elsewhere (Kataoka et al. 2000a).

If the detection of both positive and negative lags are real, it implies that cooling is sometimes but not always the dominant process, and that acceleration is also important. The (familiar) soft lag can be explained by the fact that  $\tau_{cool}$  is shorter for higher energy electrons ( $\tau_{cool} \propto (1 + u_{soft}/u_B)^{-1} B^{-2} \gamma^{-1} \delta^{-1}$ , where  $u_B$  and  $u_{soft}$  are the energy densities of the magnetic field and the soft photons to be upscattered, respectively. When high energy electrons are injected into the emitting region and then cool radiatively, the soft photons lag the hard, by a time roughly equal to the cooling time at the soft energy (Takahashi et al. 1996). Conversely, the suggested hard lag can be explained, at least qualitatively, if the acceleration time is sufficiently long,  $\tau_{acc} \sim \tau_{cool}$ , electrons appear first at low energies and gradually build up at higher energies (Georganopoulos & Marscher 1998).

The relatively small lags, also mean  $\tau_{acc} \sim \tau_{cool}$  at X-ray energies, at least during the present observations. This is consistent with the fact that the X-ray band is emitted by the highest energy regime of the electron distribution, where  $\tau_{acc}$  is approximately balanced with  $\tau_{cool}$ . According to the shock acceleration scenario,  $\tau_{acc}$  depends on energy in an opposite sense to  $\tau_{cool}$ . It follows  $\tau_{acc}(\gamma) \propto \gamma$  for diffusive acceleration with a ‘gyro-Bohm’ process (e.g. Inoue & Takahara 1996) or  $\tau_{acc}(\gamma) \propto \gamma^{1/3}$  for fully developed Kolmogorov turbulence. Similar discussions have been made by several authors (e.g. Sambruna 1999).

At a minimum, the present observations underline the need for detailed time-dependent synchrotron models, in which the time scales for electron acceleration and injection, and for radiative cooling and escape, are free to vary (e.g., Kirk et al. 1998). Depending on the relative values of these time scales, a variety of behaviors can be expected in the light curves. The fact that plateaus are rarely seen implies the dominant time scale is similar to the light-crossing time, which in turn supports the idea that the size of the emitting region is determined by diffusion and cooling.

CMU acknowledges support from NASA grant NAG5-3313. GM acknowledges NASA grants to USRA and University of Maryland.

## REFERENCES

- Aharonian, F., et al. 1999, A&A, 349, 29
- Buckley J. H., et al. 1996, ApJ, 472, L9
- Chiaberge, M., & Ghisellini, G. 1999, MNRAS, 306, 551
- Edelson, R. A., & Krolik, J. H. 1988, ApJ, 333, 646
- Fossati, G., et al., 2000a, ApJ, in press
- Fossati, G., et al., 2000b, ApJ, in press
- Georganopoulos, M., & Marscher, A. P. 1998, ApJ, 506, L11
- Ghisellini, G., et al. 1998, MNRAS, 301, 451
- Hayashida, K., et al. 1998, ApJ, 500, 642
- Inoue, S., & Takahara, F. 1996, ApJ, 463, 555
- Kataoka, J., et al. 2000a, submitted
- Kataoka, J. 2000b, PhD Thesis, Univ. of Tokyo.
- Kataoka, J., et al. 2000c, ApJ, 528, 243
- Kirk, J. G., Rieger, F.M., & Mastichiadis, A. 1998, ApJ, 332, 452
- Kubo, H. et al. 1998, ApJ, 504, 693
- Macomb, D., et al. 1995, ApJ, 449, L99 (erratum 459, L111 [1996])
- Maraschi, L., et al. 1999, ApJ, 526, L81
- Sambruna, R. M. 1999, astro-ph/9912060
- Takahashi, T., Madejski, G., and Kubo, H. 1999, APh, 11, 177 (astro-ph/9903099)
- Takahashi, T., et al. 1996, ApJ, 470, L89
- Teräsranta H., et al., 1998, A&A Supp., 132, 305
- Ulrich, M. -H., Maraschi, L., & Urry, C. M. 1997, ARAA, 35, 445
- Urry, C. M., & Padovani, P. 1995, PASP, 107, 803
- Wagner, S. J. 1996, A&A Supp., 120, 495
- Wagner, S. J., et al. 1996, AJ, 111, 2187



Wagner, S. J., & Witzel, A. 1995, ARAA 33, 163

Fig. 1.— Multiwavelength light curve for the 1998 campaign for Mkn 421. The *ASCA* light curve is extracted by integrating photons within  $1.0'$  and  $2.6'$  for SIS0 and SIS1, respectively. The small radius for the SIS0 is to avoid the saturation due to the limit of telemetry. The *EUVE* count rate is extracted from one orbit in a  $12'$  aperture. The Whipple data are limited to be that taken at elevations  $> 55$  degrees, so that the energy threshold and sensitivity is similar for all runs. Each light curve is normalized to its mean intensity. For TeV data, light curves from three telescopes are first converted to Crab unit and then the combined light curve is normalized.

Fig. 2.— Multi-frequency SED of Mrk 421 during the 1998 campaign. Filled symbols are simultaneous data obtained during the campaign. Different symbols correspond to different regions defined in Fig.1. Flux in 22 GHz, 37 GHz and R-band are averaged values during the campaign. The box from  $0.07 - 0.2$  keV is the range of the flux observed by the *EUVE* during the campaign. Butterfly-shape boxes in TeV regions are from simultaneous HEGRA observations. The EGRET data are from Macomb et al. (1996) and other data are from the NED data base. Lines are predictions by an one-zone SSC model for the steady state emission (Kataoka 2000). We fixed  $B$  to be 0.13 Gauss and  $\delta$  to be 14.  $\gamma_{max}$  and electron normalization ( $\text{cm}^{-3} \text{ s}^{-1}$ ) are  $(1.6 \times 10^5, 7.4 \times 10^{-6})$  for the quiescent state and  $(2.5 \times 10^5, 5 \times 10^{-5})$  for the high state.

Fig. 3.— Detailed time history of Mrk 421 emission obtained from *ASCA*. (a) Normalized count rate in  $0.5 - 1.0$  keV, (b)  $3 - 7.5$  keV band, (c) hardness ratio of count rates, defined as  $(3 - 7.5 \text{ keV}) / (0.5 - 1.0 \text{ keV})$ , and (d) Time lag of photons of  $0.5 - 1$  keV band from  $3 - 7.5$  keV band calculated from the DCF. Horizontal bar of each point shows the coverage of data used in the calculation.

Fig. 4.— Structure function derived from the light curve in the  $2 - 7.5$  keV and  $0.5 - 2$  keV bands with *ASCA* (circles) and in  $0.07 - 0.2$  keV band with *EUVE* (filled triangles). Open triangles show the results from ASM data in  $2 - 10$  keV band accumulated from 1996 January 6 to 1999 July 29, but normalized to the mean count rate during the present campaign.







